

(Concentrations of solutions)

Q1 a). (2018)


A student titrated 25.0 cm³ portions of dilute sulfuric acid with a 0.105 mol/dm³ sodium hydroxide solution.

Table 4 shows the student's results.

Table 4

	Titration 1	Titration 2	Titration 3	Titration 4	Titration 5
Volume of sodium hydroxide solution in cm ³	23.50	21.10	22.10	22.15	22.15

The equation for the reaction is:



Calculate the concentration of the sulfuric acid in mol/dm³

Use only the student's concordant results.

Concordant results are those within 0.10 cm³ of each other.

[5 marks]

Concentration of sulfuric acid = _____ mol/dm³

Q1. b). (2018)

Explain why the student should use a pipette to measure the dilute sulfuric acid and a burette to measure the sodium hydroxide solution.

[2 marks]

Q2. (2018)

Calculate the mass of sodium hydroxide in 30.0 cm^3 of a 0.105 mol/dm^3 solution.

Relative formula mass (M_r): $\text{NaOH} = 40$

[2 marks]


Mass of sodium hydroxide = _____ g

Q3. a). (2019)

A student investigates the volume of sodium hydroxide solution that reacts with 25.0 cm^3 of dilute sulfuric acid.

Figure 2 shows the apparatus the student uses.

Figure 2



Use **Figure 2** to answer Questions **02.3** and **02.4**

Name apparatus **A**.

[1 mark]

Q3. b). (2019)

What is the reading on apparatus **A**?

[1 mark]

_____ cm^3

Q4. (2019)

The higher the concentration of a sample of dilute sulfuric acid, the greater the volume of sodium hydroxide needed to neutralise the acid.

The student tested two samples of dilute sulfuric acid, **P** and **Q**.

Describe how the student could use titrations to find which sample, **P** or **Q**, is more concentrated.

[6 marks]


Q5. (2019)

The student repeated the investigation.

The student used solutions that had different concentrations from the first investigation.

The student found that 15.5 cm^3 of 0.500 mol/dm^3 dilute sulfuric acid completely reacted with 25.0 cm^3 of potassium hydroxide solution.

The equation for the reaction is:



Calculate the concentration of the potassium hydroxide solution in mol/dm^3 and in g/dm^3

Relative atomic masses (A_r): H = 1 O = 16 K = 39

[6 marks]

Concentration in mol/dm^3 = _____ mol/dm^3

Concentration in g/dm^3 = _____ g/dm^3

This is part of the method the student used for the titration.

1. Measure 25.0 cm^3 of the sodium hydroxide solution into a conical flask using a pipette.
2. Add a few drops of indicator to the flask.
3. Fill a burette with citric acid solution.

Describe how the student would complete the titration.

[3 marks]

Q7. (2020)

Give **two** reasons why a burette is used for the citric acid solution.

[2 marks]


1

2

Q8. (2020)

13.3 cm³ of 0.0500 mol/dm³ citric acid solution was needed to neutralise 25.0 cm³ of sodium hydroxide solution.

The equation for the reaction is:



Calculate the concentration of the sodium hydroxide solution in mol/dm³

[3 marks]

Concentration = _____ mol/dm³

Q9. (2021)

A student investigated the concentration of a solution of sodium hydroxide by titration with a 0.0480 mol/dm^3 ethanedioic acid solution.

This is the method used.

1. Measure 25.0 cm^3 of the sodium hydroxide solution into a conical flask using a 25.0 cm^3 pipette.
2. Add two drops of indicator to the sodium hydroxide solution.
3. Fill a burette with the 0.0480 mol/dm^3 ethanedioic acid solution to the 0.00 cm^3 mark.
4. Add the ethanedioic acid solution to the sodium hydroxide solution until the indicator changes colour.
5. Read the burette to find the volume of the ethanedioic acid solution used.

Suggest **two** improvements to the method that would increase the accuracy of the result.

[2 marks]

1 _____

2 _____

Ethanedioic acid is a solid at room temperature.

Calculate the mass of ethanedioic acid ($\text{H}_2\text{C}_2\text{O}_4$) needed to make 250 cm^3 of a solution with concentration 0.0480 mol/dm^3

Relative formula mass (M_r): $\text{H}_2\text{C}_2\text{O}_4 = 90$


[2 marks]

Mass = _____ g

Q11. (2021)

The student found that 25.0 cm^3 of the sodium hydroxide solution was neutralised by 15.00 cm^3 of the 0.0480 mol/dm^3 ethanedioic acid solution.

The equation for the reaction is:



Calculate the concentration of the sodium hydroxide solution in mol/dm^3

[3 marks]

Concentration = _____ mol/dm^3

Q12. a). (2022)

A student titrated 25.00 cm^3 of hydrochloric acid with 0.100 mol/dm^3 barium hydroxide solution.

Table 2 shows the results.

Table 2 shows the results.

Table 2

Titration number	1	2	3	4	5
Volume of barium hydroxide solution used in cm^3	23.90	23.45	23.55	23.55	23.45


The student calculated the volume of barium hydroxide solution to be used in the titration calculation as 23.50 cm^3 .

Explain why the student used a volume of 23.50 cm^3 of barium hydroxide solution in the titration calculation.

[2 marks]

25.00 cm³ of the hydrochloric acid reacted with 23.50 cm³ of the 0.100 mol/dm³ barium hydroxide solution.

The equation for the reaction is:




Calculate the concentration of the hydrochloric acid in mol/dm³.

[4 marks]

Concentration of the hydrochloric acid = _____ mol/dm³

Another student titrated sulfuric acid with barium hydroxide solution.


The equation for the reaction is:



The student measured the electrical conductivity of the mixture during the titration.

The better a conductor, the higher the electrical conductivity value.

Figure 10 shows the results.



Explain why the electrical conductivity of the mixture was zero when the sulfuric acid had just been neutralised.

Use the equation for the reaction.

Refer to ions in your answer.

[3 marks]

Q12. d). (2022)

The student then added a further 10 cm^3 of barium hydroxide solution.

The electrical conductivity of the mixture increased.

Give **one** reason why.

[1 mark]

Q13. (2023)

The concentration of an acid can be measured in mol/dm^3 .

Which combination of changes **increases** the concentration of an acid?

[1 mark]

Tick (✓) **one** box.

The mass of acid dissolved is halved and the volume of the solution is halved.

☐

The mass of acid dissolved is halved and the volume of the solution is doubled.

☐

The mass of acid dissolved is doubled and the volume of the solution is halved.

☐

The mass of acid dissolved is doubled and the volume of the solution is doubled.

☐

Q14. (2023)

The concentration of an acid can be determined by titration.

An indicator is added to an alkali in a flask.

Name an indicator that can be used in this titration.

Give the colour change of the indicator when acid from a burette is added to the alkali in the flask.

[2 marks]


Name of indicator _____

Colour change from _____ to _____

Q15. (2023)

A student does a titration using sodium carbonate solution and nitric acid.

The equation for the reaction is:



25.0 cm³ of 0.124 mol/dm³ sodium carbonate solution is neutralised by 23.6 cm³ of nitric acid.

Calculate the concentration of the nitric acid.

Give your answer to 3 significant figures.

You should calculate:

- the number of moles of sodium carbonate in 25.0 cm³ of the solution
- the number of moles of nitric acid in 23.6 cm³ of the nitric acid
- the concentration of the nitric acid in mol/dm³.

[5 marks]

Concentration (3 significant figures) = _____ mol/dm³

A student does a titration to find the volume of acid needed to neutralise an alkali.

The student fills a burette with the acid.

Give **three** more steps the student must do before adding the acid to the alkali from the burette.

You should name any equipment used.

[3 marks]

1 _____


2 _____

3 _____

Q17. (2024)

The student titrated a solution containing 0.0045 moles of sodium hydroxide with 0.15 mol/dm³ hydrochloric acid.

The equation for the reaction is:



Calculate the volume of hydrochloric acid in cm³ needed in the titration.

[2 marks]

Volume of acid = _____ cm³