

(Concentrations of solutions)

Q1. (2018)

$$(\text{volume} =) \frac{250}{1000} \text{ or } \frac{1}{4}$$

$$\text{or } 0.25 \text{ (dm}^3\text{)}$$

or

$$(\text{mass per cm}^3 =) \frac{40}{1000} \text{ (g)}$$

$$\text{or } 0.04 \text{ (g)}$$

$$\left(\frac{250}{1000} \times 40 = \right) 10 \text{ (g)}$$

Q2.a) (2020)

$$\frac{0.06}{25(.0)} \times 100$$

$$= 0.24 \text{ (\%)}$$

Q2.b) (2020)

(pipette) measures volume more accurately

or

(pipette has a) smaller
(percentage) uncertainty

Q3.a) (2023)

trial 3

Q3.b) (2023)

any **one** from:

- (hydrochloric acid) not added drop by drop
- did not swirl
- did not rinse apparatus (after previous trial)
- did not use a white tile
- misread pipette / burette

Q3.c) (2023)

$$\left(25.0 \text{ cm}^3 = \frac{25.0}{1000} = \right) 0.025 \text{ (dm}^3\text{)}$$

$$\text{(mass =) } 0.025 \times 4.00$$

$$= 0.1 \text{ (g)}$$

alternative approach:

$$\left(\text{concentration} = \frac{4.00}{1000} = \right) 0.004 \text{ (g/cm}^3\text{) (1)}$$

$$\text{(mass =) } 0.004 \times 25.0 \text{ (1)}$$

$$= 0.1 \text{ (g) (1)}$$