

(Extraction of metals (carbon and electrolysis))

Q1. a). (2018)

chlorine is toxic

Q1. b). (2018)

any **one** from:

- very exothermic reaction
- produces a corrosive solution
- produces hydrogen, which is explosive / flammable

Q1. c). (2018)

argon is unreactive / inert

oxygen (from air) would react
with sodium / titanium

or

water vapour (from air) would
react with sodium / titanium

Q1. d). (2018)

metal chlorides are usually ionic

(so)(metal chlorides) are solid at room temperature

or

(so)(metal chlorides) have high melting points

(because) they have strong (electrostatic) forces between the ions

or

(but) must be a small molecule or covalent

Q1. e). (2018)

sodium (atoms) lose electrons

Q1. f). (2018)



Q1. g). (2018)

(M_r of TiCl_4 =) 190

$$(\text{moles Na}) = \frac{20\,000}{23} =) \quad 870 \text{ (mol)}$$

$$\left(\text{moles TiCl}_4 = \frac{40\,000}{190} = \right) \quad 211 \text{ (mol)}$$

either

(sodium is in excess because)
870 mol Na is more than the
844 mol needed

or

(because) 211 mol TiCl_4 is less
than the 217.5 mol needed

Q1. h). (2018)

$$(\text{actual mass} =) \frac{92.3}{100} \times 13.5$$

or

$$(\text{actual mass} =) \\ 0.923 \times 13.5$$

$$= 12.5 \text{ (kg)}$$

Q2. (2019)

mixture has a lower melting
point (than aluminium oxide)

(so) less energy needed

Q3. (2019)

the electrode reacts with oxygen

the electrode is carbon /
graphite

(so) carbon dioxide is produced

Q4. a). (2021)

$$(3 \times M_r \text{ H}_2\text{O} = 3 \times (2 + 16) =) 54$$

$$(A_r \text{ R} = 150 - 54 =) 96$$

Q4. b). (2021)

(R =) molybdenum / Mo

Q5. (2021)

- carbon and iron are the cheapest reactants
- hydrogen is the most expensive reactant
- separating solid products is expensive
- separating solid products is time consuming
- in method 1, tungsten needs to be separated from tungsten carbide
- in method 1, some tungsten is lost as tungsten carbide
- in method 1, the carbon dioxide produced will escape
- in method 2, the water vapour produced will escape
- in method 2, no separation of solids is needed
- in method 3, tungsten needs to be separated from iron oxide

Q6. (2022)

both products are solid

Q7. (2023)

(substance reduced) Fe_2O_3

(reason)

(Fe_2O_3) loses oxygen