

Electrolysis of aqueous solutions (required practical)

Q1. (2019)

05.4	any two from:		2	AO3 5.4.3.4
	<ul style="list-style-type: none"> concentration / volume of solution was different impurities in solution error in timing copper falls off (electrode) copper removed when drying electrode electrode not dry (when weighed) voltage / current was different 	allow copper at bottom of beaker ignore power supply ignore recorded mass inaccurately		

Q2. (2019)

05.5		an incorrect answer for one step does not prevent allocation of marks for subsequent steps		AO2 5.4.3.4
	reading of mass at stated time	allow tolerance of $\pm \frac{1}{2}$ small square eg at 30 minutes value is 5.4 (mg)	1	
	factor from time to 24 hours	eg 5.4×48 ($= \frac{24 \text{ hours}}{30 \text{ minutes}}$) allow correct calculation using incorrectly read value for mass at time quoted	1	
	correct evaluation	eg = 259 (mg)	1	
	alternative approach:			
	calculates the gradient (1)	eg $(1.8 \div 10) = 0.18$		
	gradient $\times$ time in minutes in 24 hours (1)	eg $0.18 \times 24 \times 60$ or eg 0.18×1440 allow correct use of incorrectly determined gradient		
	correct evaluation (1)	eg = 259 (mg)		

Q3. (2021)

05.1	reduction	ignore electrolysis	1	AO2
	(as H ⁺ ions) gain electrons		1	AO1
				5.4.1.4 5.4.3.1 5.4.3.4 RPA 9

Q4. (2021)

05.2	$2\text{Cl}^- \rightarrow \text{Cl}_2 + 2\text{e}^-$	allow $2\text{Cl}^- - 2\text{e}^- \rightarrow \text{Cl}_2$ ignore state symbols allow 1 mark for $\text{Cl}_2 + \text{e}^-$ allow 1 mark for – e^- (on lhs) and Cl_2 (on rhs)	2	AO2 5.4.1.4 5.4.3.1 5.4.3.4 RPA9
------	--	--	---	--

Q5. (2021)

05.3	water molecules		1	AO1 5.4.1.4
	break down to produce OH^- ions	allow dissociate to produce OH^- ions	1	5.4.3.1 5.4.3.4 5.4.3.5 RPA9
	(which are) attracted to the positive electrode		1	
	(where OH^- ions are) oxidised or (where OH^- ions) lose electrons	ignore discharged ignore oxygen is produced as no halide is present	1	

Q6. (2022)

A)

02.1	K_2SO_4		1	AO2 5.1.1.1 5.4.3.4 RPA9
------	-------------------------	--	---	-----------------------------------

B)

02.2	(volume of hydrogen) 30 (cm^3) and (volume of oxygen) 15 (cm^3)		1	AO2 5.4.3.4 RPA9
------	--	--	---	------------------------

C)

02.3	(because) the ratio of volume of hydrogen : oxygen is 2 : 1		1	AO3 5.4.3.4 RPA9
	(and this is the) same as the ratio of hydrogen (atoms) : oxygen (atoms) in (formula of) H_2O		1	
	OR (because) the ratio of volume of hydrogen : oxygen is not 2 : 1 (1) (and this is) different to the ratio of hydrogen (atoms) : oxygen (atoms) in (formula of) H_2O (1)	must relate to the volumes given in question 02.2		

D)

02.4	$9 \pm 3 \text{ cm}^3$		1	AO2 5.3.1.4
------	------------------------	--	---	----------------

Q7. (2023)

05.3	carbon / graphite	allow platinum / gold / silver	1	AO3 5.4.3.1 5.4.3.4 RPA9
------	-------------------	--------------------------------	---	-----------------------------------

Q8. (2023)

05.5	water (molecules) break down to produce (equal numbers of) H^+ and OH^- (ions)		1	AO3
	(so) the hydrogen / H^+ ions are attracted to the negative electrode (where) hydrogen ions are discharged / reduced		1	AO1
	(because) hydrogen is less reactive than sodium		1	AO3
	(so there is) a decrease in (the relative number of) hydrogen / H^+ ions		1	AO3
	or			5.4.2.4 5.4.3.1 5.4.3.4 5.4.3.5 RPA9
	(so there is) an increase in the relative number of hydroxide / OH^- ions			