CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the October/November 2013 series

0620 CHEMISTRY

0620/31 Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

1 **(a)** uranium / plutonium / thorium [1]

(b) graphite / carbon [1]

(c) platinum / titanium / mercury / gold [1]
NOT: carbon / graphite

(d) helium [1]

(e) nitrogen / phosphorus [1]

(f) argon [1]
ACCEPT: any ion 2 + 8 + 8 e.g. K^+ etc.

(g) tellurium [1]
ACCEPT: correct symbol

[Total: 7]

2 **(a)** Any three of:
iron is harder
iron has higher density
ACCEPT: heavier **or** potassium lighter
iron has higher mp **or** bp
iron has higher tensile strength **or** stronger
iron has magnetic properties [3]
NOTE: has to be comparison, e.g. iron is hard (0) but iron is harder (1)
NOT: appearance e.g. shiny
ACCEPT: comparative statements relating to potassium

(b) potassium hydrogen (1) and potassium hydroxide (1)
zinc hydrogen (1) and zinc oxide (1)
copper no reaction (1) [5]

[Total: 8]

© Cambridge International Examinations 2013

3 (a) (i) fractional distillation [1]
(liquid) air [1]

(ii) cracking / heat in presence of catalyst [1]
of alkane / petroleum [1]
to give an alkene and hydrogen [1]

OR: electrolysis (1)
named electrolyte (1)
hydrogen at cathode (1)

OR: from methane (1)
react water / steam (1)
heat catalyst (1)
only **ACCEPT:** water with methane **or** electrolysis

(b) (i) the pair with both graphs correct is C [1]
NOTE: mark **(b)(ii)** independent of **(b)(i)**

(ii) high pressure favours side with lower volume / fewer moles [1]
this is RHS / product / ammonia [1]
$\%NH_3$ / yield increases as pressure increases [1]

the forward reaction is exothermic [1]
exothermic reactions favoured by low temperatures [1]
$\%NH_3$ / yield decreases as temperature increases [1]
ACCEPT: reverse arguments

(iii) increases reaction rate [1]
ACCEPT: reduces activation energy [1]
OR: decreases the amount of energy particles need to react
OR: economic rate at lower temperature so higher yield

[Total: 14]

4 (a) (i) (mass at t =0) – (mass at t = 5) [1]
NOTE: must have mass at t = 5 not final mass

(ii) fastest at origin
slowing down between origin and flat section gradient = 0
where gradrient = 0
three of above in approximately the correct positions [2]

(iii) 3 correct comments about gradient = [2]
2 correct comments about gradient = [1]
1 correct comment about gradient = [0] [2]

(b) start at origin and smaller gradient [1]
same final mass just approximate rather than exact [1]

© Cambridge International Examinations 2013

(c) (i) smaller surface area [1]
lower collision rate [1]

(ii) molecules have more energy [1]
collide more frequently / more molecules have enough energy to react [1]

(d) number of moles of HCl in 40 cm^3 of hydrochloric acid,
concentration 2.0 mol / dm^3 = 0.04 × 2.0 = 0.08 [1]
maximum number of moles of CO_2 formed = 0.04 [1]
mass of one mole of CO_2 = 44 g [1]
maximum mass of CO_2 lost = 0.04 × 44 = 1.76 g [1]

[Total: 15]

5 (a) (i) have same molecular formula / both are C_5H_{12} [1]
they have different structural formulae / different structures [1]

(ii) CH_3-CH_2-CH=CH-CH_3 / any other correct isomer [1]

(b) (i) CH_2-(Br)-CH_2Br [1]
NOT: $C_2H_4Br_2$
dibromoethane [1]
NOTE: numbers not required but if given must be 1, 2

(ii) CH_3-CH_2-CH_3 [1]
NOT: C_3H_8
propane [1]

(iii) CH_3-CH_2-CH_2-CH_2-OH / CH_3-CH_2-CH(OH)-CH_3 [1]
butanol [1]
numbers not required but if given must be correct and match formula

(c) (i) CH_3-CH=CH-CH_2-CH_3 [1]
CH_3-CH=CH-CH_3 [1]

(ii) pink / purple [1]
colourless [1]
NOT: clear

(d) -CH_2-CH(CN)-CH_2-CH(CN)-
correct repeat unit CH_2-CH(CN) [1]
COND: at least 2 units in diagram [1]
continuation [1]

[Total:16]

© Cambridge International Examinations 2013

6 (a) (i) (attractive force between) positive ions [1]
and (negative) electrons [1]
opposite charges attract ONLY [1]
electrostatic attraction ONLY [1]

(ii) lattice / rows / layers of lead ions / cations / positive ions [1]
NOT: atoms / protons / nuclei
can slide past each other / the bonds are non-directional [1]

(b) (i) anhydrous cobalt chloride becomes hydrated [1]
ACCEPT: hydrous

(ii) carbon dioxide is acidic [1]
sodium hydroxide and calcium oxide are bases / alkalis [1]

(iii) Any two of:
water, calcium carbonate and sodium carbonate [2]
ACCEPT: sodium bicarbonate

(c) number of moles of CO_2 formed = 2.112 / 44 = 0.048 [1]
number of moles of H_2O formed = 0.432 / 18 = 0.024 [1]

x = 2 and y = 1 **NOT:** ecf from this line

formula is $2PbCO_3.Pb(OH)_2$ / $Pb(OH)_2.\ 2PbCO_3$ [1]

[Total:12]

7 (a) (i) hydrogen (atoms) replaced by (atoms) of a different element e.g. chlorine [1]
NOT: substitute

(ii) light required [1]

(b) exothermic reaction gives out energy [1]
endothermic reaction absorbs
takes in energy [1]

(c)

bonds broken	energy
C-H	+412
C*l*-C*l*	+242
total energy	+654

[1]

bonds formed	energy
C-C*l*	–338
H-C*l*	–431
total energy	–769

[1]
energy change –115 [1]
negative sign indicates exothermic [1]

[Total: 8]

© Cambridge International Examinations 2013