

## AQA GCSE Physics — Physics

### Topic 6

# Waves


Specification 8463 · AQA

## Free edition


**Revision notes for one topic, free to download.**

These are the summary notes only. The full edition of this booklet also carries the worked examples, the required practicals and the formula sheets, with every topic in one place.

## Colour Codes



**Definitions**



**Diagrams**



**Notes and warnings**



**Sub-topic headings**

This booklet is a study aid written from the published specification.  
It is not affiliated with, endorsed by or approved by AQA.

**Free to download and share for personal study.**  
**Please do not sell or republish it as your own.**  
**For enquiries, please email [pcfgyrdi@gmail.com](mailto:pcfgyrdi@gmail.com)**

### Define Transverse wave

- oscillations perpendicular to the direction of energy transfer.

### Define Compression

- region of a longitudinal wave where particles are pushed together.

### Define Amplitude

- the maximum displacement of a point on a wave from its undisturbed position.

### Define Frequency

- the number of waves passing a point each second, in hertz (Hz).

### Define Wave speed

- the speed at which the energy is transferred (or the wave moves) through the medium.

### Define Specular reflection

- reflection from a smooth surface in a single direction.

### Define Diffuse reflection

- reflection from a rough surface, causing scattering.

### Define P-wave

- longitudinal seismic wave; travels through solids and liquids.

### Define Principal focus

- the point where a convex lens brings parallel rays to a focus.

### Define Longitudinal wave

- oscillations parallel to the direction of energy transfer.

### Define Rarefaction

- region of a longitudinal wave where particles are spread apart.

### Define Wavelength

- the distance from a point on one wave to the equivalent point on the adjacent wave.

### Define Period

- the time for one complete wave to pass a point, in seconds (s).

### Define Reflection

- a wave bouncing back at a boundary between two materials.

### Define Refraction

- a wave changing direction at a boundary because its velocity changes.

### Define Ultrasound

- sound of frequency above the upper limit of human hearing.

### Define S-wave

- transverse seismic wave; cannot travel through a liquid.

### Define Focal length

- the distance from the lens to the principal focus.

### Define Radiation dose

- a measure, in sieverts, of the risk of harm from exposure to radiation.

### Define Perfect black body

- an object that absorbs all radiation incident on it, reflecting and transmitting none.

## 6.1 Waves in air, fluids and solids

### 6.1.1 Transverse and longitudinal waves

Waves may be either **transverse** or **longitudinal**. Both transfer energy from a source to an absorber.


|             | Transverse                                           | Longitudinal                                |
|-------------|------------------------------------------------------|---------------------------------------------|
| Oscillation | Perpendicular to energy transfer                     | Parallel to energy transfer                 |
| Features    | Peaks and troughs                                    | <b>Compressions</b> and <b>rarefactions</b> |
| Examples    | Ripples on water, all electromagnetic waves, S-waves | Sound in air, P-waves                       |

### Evidence that the wave travels, not the medium

- A cork on a pond bobs **up and down** as ripples pass but is not carried to the far bank.
- A loudspeaker makes air particles vibrate about fixed positions. If the air itself travelled to your ear there would be a wind, and the room would empty of air.

## Transverse and longitudinal waves compared

### Transverse wave — a wave sent along a rope



The oscillations are at right angles ( $90^\circ$ ) to the direction of energy transfer.  
Examples: ripples on water, all electromagnetic waves, S-waves.

### Longitudinal wave — a wave sent along a slinky spring

compression — coils pushed together

rarefaction — coils spread apart



Examples: sound waves travelling through air, P-waves (seismic waves).

### Waves transfer energy (and information) — they do not transfer matter

float bobs up and down



A float resting on water ripples, or a piece of paper in front of a loudspeaker, just oscillates about a fixed point — it is not carried along with the wave. This is evidence that it is the wave (the disturbance) that travels, and not the water or the air itself.

#### ANALOGY

**Rope vs slinky.** Flick a **rope** and a hump runs along it — your hand moves at right angles to the hump's travel: transverse. Push and pull a **slinky** and a squashed patch runs along it — your hand moves along the line the wave travels: longitudinal. Both end up where they started; only energy moves along.

#### WATCH OUT


**"Waves carry the water or air along with them."** Waves transfer **energy**, not **matter**. Particles of the medium oscillate about a fixed undisturbed position and stay there. Answer as "the wave transfers energy from the source to an absorber; the particles only vibrate".

### 6.1.2 Properties of waves

- **Amplitude** is measured from the middle (undisturbed) line to a peak — **not** peak to trough.
- **Wavelength** ( $\lambda$ ) is peak to peak, or compression to compression.
- **Wave speed** is the speed at which energy is transferred through the medium.
- You must identify amplitude and wavelength from given diagrams. On a displacement–distance graph the x-axis gives the wavelength; on a displacement–time graph one wave gives the **period**.

### The parts of a wave: wavelength, amplitude, peak and trough

#### Parts of a wave



#### The equations

wave speed = frequency  $\times$  wavelength

$$v = f \lambda$$

wave speed,  $v$ , in metres per second, m/s

frequency,  $f$ , in hertz, Hz

wavelength,  $\lambda$ , in metres, m

period =  $1 \div$  frequency

$$T = 1 / f$$

period,  $T$ , in seconds, s

frequency,  $f$ , in hertz, Hz

frequency = waves passing a point each second

### Measuring the speed of sound waves in air


1. Connect two microphones a measured distance apart (about 1 m) to an oscilloscope.
2. Make a sharp sound beyond the first microphone; measure the **time delay** between the two traces.
3. speed = distance  $\div$  time delay. Repeat and mean (about 330–340 m/s).
4. Or clap a measured distance from a large wall and time 20 echoes; total distance =  $2 \times \text{distance} \times 20$ .

**Measuring the speed of ripples on a water surface.** Use a **ripple tank**, as in required practical 8.

**PHYSICS ONLY** Passing from one medium into another, the **frequency stays the same** (set by the source), so from  $v = f \lambda$  a higher velocity means a proportionately longer wavelength.

### Measuring frequency, wavelength and speed in a ripple tank

#### Required practical 8 — measuring waves in a ripple tank




Frequency  $f$ : count the waves passing a point in 10 s and divide by 10.

Wavelength  $\lambda$ : measure across 10 wavelengths with the metre rule and divide by 10.

Wave speed:  $v = f \lambda$ . A stroboscope freezes the pattern, making it easier to measure.

#### Waves in a solid — a stretched string



### 6.1.3 Reflection of waves **PHYSICS ONLY**

**PHYSICS ONLY** At a boundary between two materials a wave can be:

- **reflected** — it bounces back into the first material;
- **transmitted** — it passes into the second material, usually refracting as it does;
- **absorbed** — its energy is transferred to the second material's energy stores, heating it.


**PHYSICS ONLY** Which happens depends on the two materials and on the wavelength; usually all three occur at once.

**PHYSICS ONLY** **Ray diagrams for reflection.** Draw the surface and the **incident ray**; add the **normal** (dashed, at  $90^\circ$  to the surface where the ray strikes); draw the **reflected ray** on the other side of the normal with the angle of reflection **equal** to the angle of incidence, with arrows. Angles are always measured **from the normal**.

### Specular and diffuse reflection, and refraction through a glass block

Physics only

#### Specular reflection — smooth surface



All the reflected rays travel in the same direction, so a clear image is formed.


#### Diffuse reflection — rough surface

parallel rays of light arrive




Each ray still obeys  $i = r$ , but the dashed normal points a different way at each point, so light is scattered.

#### Refraction through a glass block



Entering glass the light slows down and bends towards the normal. Leaving, it speeds up and bends away again.

#### Why it happens — wavefronts



The edge of each wavefront that reaches the glass first slows down, so the whole wave swings round and changes direction.

The normal is a construction line drawn at  $90^\circ$  to the surface, where the ray meets it.

$i$  = angle of incidence;  $r$  = angle of reflection, or angle of refraction.

6.1.4 Sound waves **PHYSICS ONLY****HT ONLY**

**PHYSICS ONLY HT ONLY** Sound waves can travel through **solids**, causing vibrations in the solid. Within the **ear**, sound waves cause the **ear drum** and other parts to vibrate, causing the sensation of sound.

**PHYSICS ONLY HT ONLY** This conversion only works over a **limited frequency range**, which restricts the limits of human hearing: normal human hearing runs from **20 Hz to 20 kHz**. Frequencies outside it do not make the parts of the ear vibrate efficiently, so are not heard.

**PHYSICS ONLY HT ONLY** Sound travels at **different speeds in different media** because the speed depends on how strongly the particles are coupled: in a solid they are close and strongly bonded, so a disturbance passes on quickly; in a gas they are far apart, so it travels slowly. Sound is fastest in solids, slowest in gases.

6.1.5 Waves for detection and exploration **PHYSICS ONLY****HT ONLY**

**PHYSICS ONLY HT ONLY** Differences in **velocity, absorption and reflection** between different types of wave in solids and liquids let us detect and explore structures hidden from direct observation.

**Ultrasound**

- **Ultrasound waves** have a frequency higher than the upper limit of hearing for humans (above 20 kHz).
- They are **partially reflected** at a boundary between two different media; the **time taken** for the reflections to reach a detector gives the distance to that boundary.
- Used for **medical imaging** (pre-natal scans, soft tissue) and **industrial imaging** (cracks inside castings).

**Echo sounding** uses **high frequency sound waves** to detect objects in deep water and measure water depth.

**Seismic waves**


- Seismic waves are produced by **earthquakes**. **P-waves** are longitudinal and travel at different speeds through solids and liquids, passing through both; **S-waves** are transverse and **cannot travel through a liquid**.
- An **S-wave shadow zone** on the far side of the Earth, where no S-waves arrive, is evidence that part of the core is **liquid**. P-waves arrive but are **refracted** as their speed changes; the shadow zones give the **size of the core**.
- **P-waves and S-waves provide evidence for the structure and size of the Earth's core** — parts of the Earth that are **not directly observable**.

## 6.2 Electromagnetic waves

## 6.2.1 Types of electromagnetic waves

- Electromagnetic waves are **transverse** waves that transfer energy from the **source** of the waves to an **absorber**. They form a **continuous spectrum**, and all types travel at the same velocity through a vacuum (space) or air —  $3 \times 10^8$  m/s.
- Grouped by **wavelength** and **frequency**, from **long to short wavelength (low to high frequency)**: radio, microwave, infrared, visible light (red to violet), ultraviolet, X-rays, gamma rays.
- Our **eyes only detect visible light**, so we detect a limited range of electromagnetic waves.
- Energy transfer examples: a microwave oven heating food; sunlight warming a pavement; an aerial absorbing radio waves to produce a current.

## The electromagnetic spectrum, from radio waves to gamma rays



| Group         | One use                            | One hazard                      |
|---------------|------------------------------------|---------------------------------|
| radio waves   | television and radio broadcasts    | very large doses heat tissue    |
| microwaves    | satellite links, cooking food      | internal heating of body tissue |
| infrared      | electric heaters, infrared cameras | skin burns                      |
| visible light | fibre optic communications         | retina damage if very bright    |
| ultraviolet   | energy efficient lamps, tanning    | skin ages early; skin cancer    |
| X-rays        | medical imaging (X-ray photos)     | ionising: mutation and cancer   |
| gamma rays    | sterilising food, treating cancer  | ionising: mutation and cancer   |

All electromagnetic waves are transverse and transfer energy from a source to an absorber. They form a continuous spectrum and all travel at the same speed through a vacuum (space) or air. Visible light runs from red (longest wavelength) to violet (shortest); our eyes detect only this narrow band.

### WATCH OUT

**Getting the spectrum the wrong way round.** Use **Rich Men In Vegas Use eXpensive Gold**. In that order **wavelength decreases, frequency increases** and the **energy carried increases** — which is why the ionising ones (UV, X-rays, gamma) are at the gamma end. What does **not** change is the **speed**: in a vacuum every part travels at the same speed, so gamma rays are not "faster" than radio waves, only higher in frequency.

### ANALOGY

**A car driving onto sand.** A car leaves the road onto soft sand at an angle: the wheel reaching the sand first slows while the other is still on tarmac, so the car **slews round**. Drive straight on (along the normal) and both wheels slow together, so it slows but carries straight on. Refraction happens for exactly that reason — the **change of speed**.

### HIGHER TIER

#### 6.2.2 Properties of electromagnetic waves 1

**HT ONLY** Different substances may **absorb, transmit, refract or reflect** electromagnetic waves in ways that **vary with wavelength** — glass transmits visible light but absorbs most ultraviolet.

**HT ONLY** Some effects, for example **refraction**, are due to the difference in velocity of the waves in different substances. When a wave meets a boundary at an angle, one side of the wavefront changes speed **before** the other, swinging it round so the direction of travel changes. Slowing down bends the ray **towards** the normal, speeding up bends it **away**; a wave arriving **along** the normal changes speed and wavelength but not direction.

**Ray diagrams for refraction.** Draw the boundary and the **normal** (dashed, at 90° to it), then the incident ray marking the **angle of incidence** from the normal, then the refracted ray bent towards or away from the normal as above.

**HT ONLY** On a **wavefront diagram**, draw wavefronts as parallel lines at right angles to the ray: in the slower medium they are **closer together** (shorter wavelength, frequency unchanged) and **tilted** relative to the first medium.

### 6.2.3 Properties of electromagnetic waves 2

**HT ONLY** Radio waves can be produced by oscillations in electrical circuits. An alternating current in a transmitting aerial makes electrons oscillate, and these oscillating charges emit radio waves of the **same frequency** as the current.

**HT ONLY** The reverse also happens: when radio waves are **absorbed** they may create an alternating current with the same frequency as the radio wave itself, so radio waves can induce oscillations in an electrical circuit.

- Changes in **atoms and the nuclei of atoms** can generate or absorb electromagnetic waves over a wide frequency range; gamma rays originate from changes in the nucleus of an atom.
- Ultraviolet waves, X-rays and gamma rays can have hazardous effects on human body tissue, depending on the **type of radiation** and the **size of the dose**.
- **Radiation dose (in sieverts)** measures the **risk of harm** from exposure of the body to radiation; 1000 millisieverts (mSv) = 1 sievert (Sv). You need not recall this unit, but may be asked to draw conclusions from given data about the risks of exposure.
- **Ultraviolet** can cause skin to **age prematurely** and increases the risk of **skin cancer**. **X-rays and gamma rays** are **ionising radiation** that can cause the **mutation of genes** and **cancer**.

### 6.2.4 Uses and applications of electromagnetic waves

| Part of spectrum | Use                                                | Hazard                                              |
|------------------|----------------------------------------------------|-----------------------------------------------------|
| Radio waves      | Television and radio                               | No significant hazard — lowest frequency and energy |
| Microwaves       | Satellite communications; cooking food             | Internal heating of body tissue                     |
| Infrared         | Electrical heaters; cooking food; infrared cameras | Skin burns                                          |
| Visible light    | Fibre optic communications                         | Retinal damage if very intense                      |
| Ultraviolet      | Energy efficient lamps; sun tanning                | Premature skin ageing; skin cancer                  |
| X-rays           | Medical imaging and treatments                     | Ionising: mutation of genes, cancer                 |
| Gamma rays       | Medical imaging and treatments                     | Ionising: mutation of genes, cancer                 |

#### NOTE

The specification only names hazards for **ultraviolet, X-rays and gamma rays**; the others are given for completeness and are not required.

### 6.2.5 Lenses **PHYSICS ONLY**

**PHYSICS ONLY** A **lens** forms an image by **refracting** light. In a **convex** lens, parallel rays are brought to a focus at the **principal focus**, a distance equal to the **focal length** from the lens.

**HT ONLY** Why each type suits its application:


- **Radio** — barely absorbed by the atmosphere, so covers wide areas.
- **Microwaves** — pass **through** the atmosphere to satellites; oven wavelengths are absorbed by water in food.
- **Infrared** — strongly absorbed by surfaces, heating them; all warm objects emit it, so cameras detect heat.
- **Visible light** — weakly absorbed by glass fibres and reflects along them, so data is carried with little loss.
- **Ultraviolet** — makes fluorescent coatings glow, wasting less energy.
- **X-rays** — pass through flesh but are absorbed by bone, giving a shadow image; **gamma rays** penetrate the body and kill cancer cells.

|               | Convex (converging)                           | Concave (diverging)                                    |
|---------------|-----------------------------------------------|--------------------------------------------------------|
| Shape         | Thicker in the middle                         | Thinner in the middle                                  |
| Parallel rays | Brought together at the principal focus       | Spread out, appearing to come from the principal focus |
| Image         | <b>Real or virtual</b>                        | <b>Always virtual</b>                                  |
| Symbol        | Line with <b>outward</b> -pointing arrowheads | Line with <b>inward</b> -pointing arrowheads           |


#### PHYSICS ONLY Drawing the ray diagrams

1. Draw the principal axis and lens symbol, marking **F** at the focal length on both sides; draw the object as an upright arrow.
2. **Ray 1:** from the top of the object, parallel to the axis, then through F (convex) or appearing to come from F (concave).
3. **Ray 2:** from the top of the object straight through the **centre** of the lens, undeviated.
4. The rays cross at the top of the image. A **real** image forms where rays actually meet: it can be projected and is inverted. A **virtual** image forms where rays only **appear** to come from, traced **back** as dashed lines: it cannot be projected and is upright.


**Convex (converging) lens — object well beyond the principal focus**



**Convex lens — object closer than the principal focus (magnifying glass)**



**Concave (diverging) lens — the image is always virtual**



**Magnification**

$$\text{magnification} = \text{image height} \div \text{object height}$$

Magnification is a ratio, so it has no units. Measure both heights in the same unit (mm or cm).

## 6.2.6 Visible light PHYSICS ONLY

**PHYSICS ONLY** Each **colour** in the visible light spectrum has its own **narrow band of wavelength and frequency** — red the longest wavelength and lowest frequency, violet the shortest and highest. Reflection from a **smooth** surface in a single direction is **specular reflection**; reflection from a **rough** surface causes scattering, called **diffuse reflection**.

**PHYSICS ONLY** The colour of an **opaque** object is determined by **which wavelengths are more strongly reflected**; the rest are absorbed. If **all** are reflected equally it appears **white**; if all are absorbed it appears **black**. A red jumper reflects red and absorbs the rest, so in pure blue light it looks black.

**PHYSICS ONLY** Objects that transmit light are either **transparent** (a clear image through them) or **translucent** (light passes but is scattered, blurring the image).

**PHYSICS ONLY** **Colour filters** work by absorbing certain wavelengths (and colour) and transmitting other wavelengths (and colour): a red filter transmits red and absorbs the rest, so white light comes out red. Red and blue filters together let almost nothing through.

## 6.3 Black body radiation PHYSICS ONLY

### 6.3.1 Emission and absorption of infrared radiation

**PHYSICS ONLY** All bodies (objects), no matter what temperature, emit and absorb infrared radiation, and the **hotter** the body the **more** infrared radiation it radiates in a given time.

**PHYSICS ONLY** A **perfect black body** absorbs **all** the radiation incident on it and does **not reflect or transmit** any. Since a good absorber is also a good emitter, a perfect black body would be the best possible emitter.

#### ANALOGY

**Why black is best at both.** Think of a surface as a doorway between the radiation outside and the energy stores inside. Matt black is a wide-open door: radiation walks straight in and nothing bounces off, so it is the best **absorber** — but an open door lets traffic both ways, so energy leaves as easily, making it the best **emitter** too. Shiny silver is a closed, mirrored door: poor at both.

### 6.3.2 Perfect black bodies and radiation

**PHYSICS ONLY** All bodies emit radiation, whatever their temperature, and the **intensity** and **wavelength distribution** of the emission **depend on the temperature** of the body: as a body gets hotter it emits more radiation overall **and** the peak moves to **shorter wavelengths** — a heating element glows dull red, then orange, then white.

**PHYSICS ONLY HT ONLY** A body at **constant temperature** is absorbing radiation at the **same rate** as it is emitting radiation; its temperature **increases** when it absorbs faster than it emits, and falls when it emits faster. Tea cools because it emits more than it absorbs; a sunlit road warms because it absorbs more, until the rates balance.

**PHYSICS ONLY HT ONLY** The temperature of the Earth depends on many factors, including:

- the **rate of absorption** of radiation — mainly short-wavelength radiation from the Sun absorbed by the surface;
- the **rate of emission** of radiation — the Earth emits longer-wavelength **infrared** back towards space;
- the **reflection of radiation into space** by clouds, ice and light-coloured surfaces before it is absorbed.

**PHYSICS ONLY HT ONLY** The surface warms by day, when absorption exceeds emission, and cools at night; the average temperature stays constant only while radiation absorbed equals radiation emitted. Atmospheric gases absorb some outgoing infrared and re-radiate part back to the surface, slowing energy loss, so if these gases increase the average temperature **rises** until balance is restored. More reflective ice or cloud lowers it.